

RESEARCH ARTICLE

The Annals of Zoology

ISSN (Print): 0003-5009

Annals of Zoology, Vol. 31; December 2015, 28-32

©All Rights Reserved Council of Natural Sciences, India

Hydrobiological Factors Related with Augmentation for Fish Production in Certain Selected Ponds of Saran

Prabhat Ranjan¹ and Rabindra Kumar²

¹ S.M.D. College, MN Jalalpur, Gopalganj (Bihar)

² P.G. Department of Zoology, Jai Prakash University, Chapra (Bihar)

Email: rounak.raj9@gmail.com

ABSTRACT

A hydro-biological study conducted in four different ponds of Chapra district, showed that the concentrations of chemical parameters like dissolved oxygen, free carbon dioxide, pH, conductivity, alkalinity, nitrate, phosphate, calcium, magnesium, copper and zinc are within the permissible levels of drinking water quality standard of WHO and ISI. However, iron content was higher in most of the ponds. A clear indirect relationship between iron concentration and euglenoids has been observed. Major phytoplankton taxa present in the ponds are Chlorophyceae, Cyanophyceae, Bacillariophyceae and Euglenophyceae. The study reveals that rural ponds can be a very good source of water for drinking, domestic use and fishery and should be conserved at any cost.

Key words: Drinking water quality, Iron, Euglenophyceae, Chlorophyceae

INTRODUCTION

The world's water resources are under pressure and must be managed for human survival. It is, therefore, necessary to have most relevant information for arriving at rational decisions that will result in the maximum benefit to most people. Accurate and reliable information on the water resource system can, therefore, be a vital aid to strategic management of the resources (Gupta and Deshpande, 2004). Ponds have been used since time immemorial as a traditional source of water supply in India. However, the water of the ponds, lakes and river are polluted mainly due to discharged waste water from residential areas, sewage outlets, solid wastes, detergents, automobile oil wastes, fishing facilities and agricultural pesticides from farmlands (Usha, *et al.*, 2006; Hasan, *et al.*, 2007). In recent years, their importance has somewhat declined due to technological advancements leading to more centralized water supply systems. There is a relationship among ecologists and micro-planners about the importance of conservation of ponds as sustainable source of water for rural communities (Park and Park, 2005).

The present study is an attempt to assess the water quality of ponds in Chapra district so that they may be sustainably exploited for multiple uses like rural water supply, fisheries and even recreation.

METHODS AND MATERIALS

The study was carried out in four different ponds of Chapra district in which 1 and 3 are household ponds while pond 2 and 4 are fishery ponds.

Water samples were collected fortnightly from February to April, 2003 from the sub-surface layer of the ponds in PVC and BOD bottles (for estimating dissolved oxygen). Hydrogen ion concentration and conductivity were measured by a pH and conductivity meter, respectively, while free carbon dioxide, dissolved oxygen and total alkalinity were estimated by standard methods (Ramesh and Anbu, 1996). Nitrate content was measured by Ion Meter with Ion Selective electrode (Eutech, Germany) while phosphate content was estimated by Ammonium Molybdate-Stannous Chloride method (Michael, 1984).

Ca, Mg, Fe, Cu and Zn contents were estimated (Gupta, 1996) by Perkin-Elmer 2380 flame atomic absorption spectro-photometer. The readings were checked with those of standard solutions and procedural blanks, acid washed glassware, analytical grade reagents and double distilled de-ionized water were used to minimize contamination errors. The detection limits for Ca, Mg, Fe, Cu and Zn were 1.0, 0.1, 3.0, 1.0 and 0.8 g l^{-1} respectively. Quantitative sampling of plankton was done by filtering a known volume of water through plankton net. After fixation in 6 percent formalin and pre-concentration, cell counts were made in a sedgewick rafter (Michael, 1984). Statistical analysis was done by using window based statistical package SYSTAT 10.

RESULTS AND DISCUSSIONS

Quality of an aquatic ecosystem is dependent on the physicochemical qualities of water as also on the biological diversity of the system (Tiwari and Chauhan, 2006). Cairns and Dickson (1971) stated that analysis of biological materials along with chemical factors of water forms a valid method of water quality assessment.

Table 1: Variation of chemical parameters in ponds 1 to 4

| Var iables | Pond1 | Pond2 | Pond3 | Pond4 |
|----------------------|--------------|--------------|---------------|---------------|
| DO | 6.37 (1.01) | 8.19 (1.07) | 6.55 (2.40) | 5.91 (2.38) |
| Free CO ₂ | 12.47 (3.36) | 13.27 (4.68) | 12.53 (0.53) | 23.47 (16.66) |
| TA | 20.00 (3.60) | 11.00 (3.6) | 17.00 (6.10) | 47.87 (56.23) |
| pH | 7.40 (0.34) | 7.47 (0.25) | 7.24 (0.14) | 7.5 (0.21) |
| Cnd. | 123.8 (8.26) | 29.63 (1.7) | 66.4 (9.93) | 114.3 (36.96) |
| TDS | 56.30 (3.79) | 14.00 (1.0) | 28.90 (7.10) | 52.67 (16.56) |
| Nitrate | 0.720 (0.38) | 0.210 (0.11) | 0.170 (1.15) | 0.38 (0.32) |
| Phosphate | 0.850 (0.01) | 0.00 | 0.49 (0.04) | 2.56 (0.25) |
| Calcium | 0.013 (0.01) | 0.03 (0.01) | 0.70 (0.02) | 0.14 (0.001) |
| Magnesium | 5.15 (0.05) | 2.05 (0.05) | 04.70 (0.08) | 3.37 (0.01) |
| Iron | 1.13 (0.01) | 0.71 (0.045) | 01.78 (0.02) | 0.49 (0.01) |
| Copper | 0.081 (0.01) | 0.07 (0.01) | 0.110 (0.025) | 0.09 (0.02) |
| Zinc | 0.820 (0.03) | 0.39 (0.02) | 0.300 (0.01) | 0.30 (0.01) |

Table 1 depicts the chemical variables of the water of four different ponds selected for the present study. Variations in percentage occurrence of major phytoplankton taxa in ponds 1 to 4 are shown in Fig. 1. Major taxa present are Chlorophyceae, Cyanophyceae, Bacillariophyceae and Euglenophyceae. The planktonic algae are particularly sensitive to chemical changes and myriad environmental conditions promote the development of the algal spores present in the sediment (Rodhe, 1948). The pond is used only for washing and bathing without entry of sewage and domestic waste. Hence the two groups, Cyanophyceae and Euglenophyceae which are generally seen to appear near sewage outfall (Pandit, 2002) are not encountered in the pond.

The solubility and availability of nutrients is affected by oxygen content of water and therefore the productivity of aquatic ecosystems (Wetzel, 1984). The highest dissolved oxygen value and nearly neutral pH in pond 4 can be attributed to the diversified nature of the plankton population of the pond where family, Euglenophyceae was the most dominant group followed by

Chlorophyceae: As chlorophyll-b is possessed by these two groups only besides possessing chlorophyll-a, they must have transferred an additional amount of light energy to chlorophyll-a for primary photo chemical reactions (Wetzel, 1984). The lowest dissolved oxygen value in pond 3 associated with the highest levels of conductivity, total dissolved solids and pH, as well as high free carbon dioxide and iron values is in agreement with studies conducted at Jaisamand lake, Udaipur, Rajasthan (Sharma and Sarang, 2004) and Naukuchiyatal lake in Kumaon Himalaya (Pandey and Soni, 1993). Algal bloom in pond 3 also confirms the relationship of high pH value with phytoplankton bloom (Mitra and Bhattachariya, 1965). The chlorophytic algal bloom in this pond might

have been instrumental in using up oxygen during the decomposition of the dying bloom which in turn led to the release of the nutrients.

Table 2: Significant correlations among chemical variables

| Variables | r | p | Variables | r | p |
|-------------------------|--------|--------|------------------------|--------|---------|
| DO vs. CO ₂ | -0.365 | <0.01 | Ca vs. TA | 0.321 | < 0.02 |
| DO vs. NO ₃ | 0.561 | <0.001 | PO ₄ vs. TA | 0.350 | < 0.01 |
| Cnd.vs. PO ₄ | 0.510 | <0.001 | pH vs. NO ₃ | -0.336 | < 0.02 |
| Cnd. vs. Mg | 0.289 | <0.05 | Fe vs. Cu | 0.683 | < 0.001 |
| Ca vs. Mg | 0.350 | <0.01 | Fe vs. Zn | 0.445 | < 0.001 |

In lake ecosystems beside the input of nitrate through runoff, decomposition of nitrogenous matter and its further oxidation plays a significant role. All ponds have been found to be favorable for fish productivity as nitrate value of these ponds ranged between 0.1-3.0 mg l⁻¹ (Verma, 2002). The low range of phosphate value in all the ponds is due to the fact that at high temperature, phosphate is rapidly assimilated by plankton and microorganisms (Manna and Das, 2004).

In this study, we find that concentration of iron is beyond the permissible level of water quality standard in ponds 1 and 3 while other parameters are within the limits in all the ponds. Gobler and Cospers (1996) and Indian worker (Khan and Bhat, 2000), have recorded much higher concentration of iron in lake and pond waters and emphasized its role in inducing chrysophyte and euglenoid blooms regularly. It has been observed that it is totally absent in the ponds where iron level is quite high where as in pond 4 comparatively lower iron value is associated with moderate abundance of Euglenophyceae. From this study and previous studies, it can be said that as in these types of shallow water bodies boom and bust cycle continues with the entry of nutrients via surface runoff, concentration of iron in the pond 1, 2 and 3 was not conducive for the initiation of the growth of euglenoids while in ponds 4 with the optimum concentration of iron euglenoids started to develop and spread, thus utilizing and reducing the iron content of water. This confirms other limnological study, which showed that besides iron, calcium and magnesium also have a great role in stimulating and maintaining *Euglena* blooms (Dutta Gupta, *et al.*, 2004). This is possible because calcium increases the availability of other ions and magnesium acts as a carrier of phosphorus (Wetzel, 1984). Lake Manasbal of Kashmir valley of India which is infested by *Euglena pedunculata* also has calcium rich water (Khan and Bhat, 2000). The concentration of copper in all the ponds have been found to be very low compared to other elements and concentration of zinc also has not shown much fluctuation. Though zinc is involved in nucleic acid synthesis and participates in a variety of metabolic processes involving carbohydrates, lipids, proteins and nucleic acid (McDowell, 1992), it can be toxic also when present in excess amount as changes in blood parameters and tissue structures have been reported on exposure to zinc (Banerjee, 1998).

Correlation coefficients computed among the chemical parameters of nine ponds showed a number of significant relationships (Table 2). The classical inverse relationship between dissolved oxygen and carbon dioxide was found to be significant as dissolved oxygen is mainly regulated by photosynthetic activity of algal flora when free carbon dioxide is utilized (Wetzel, 1984). In contrast, positive relationship between dissolved oxygen and nitrate may be explained by the fact that nitrate concentrations in all the ponds were not very high, the maximum value being 2.2 mg l⁻¹. The moderate nitrate concentration is likely to have enhanced the growth of phytoplankton which in turn produced more dissolved oxygen. Significant positive correlations of conductivity with phosphate and magnesium indicate that they are the key factors governing the conductivity regimes of the ponds investigated.

Phosphate is essential for the growth of organisms and can limit primary productivity (Wetzel, 1984). Again magnesium is required by the chlorophyllous plants as it acts as a micronutrient in enzymatic trans-phosphorylation by algae. It was also shown that low available magnesium can influence the phytoplankton productivity in oligotrophic Alaskan lakes (Goldman, 1960). Calcium and magnesium are significantly correlated which can be attributed to the fact that both are integral part of plant tissue and contribute to the hardness of water (Wetzel, 1984). Further they play an important role in neutralising the excess acid produced in the system (Das, 2002). This also justifies the significant positive relationship of calcium with alkalinity. Inverse relationship of pH with nitrate as recorded in the present study agrees well with the study made in the Maheshkhali Channel, Bay of Bengal, Bangladesh (Ali, *et al.*, 1985). Iron showed significant positive correlation with copper and zinc. These are essential micronutrients for plants and many animals, required in trace amounts, and thus vital in the molecular architecture of various proteins, enzymes and vitamins (Giddings, 1973).




Fig.1: Dominant Phytoplankton groups in studied Ponds.

The present findings indicate that water quality of all the nine ponds is suitable for drinking and domestic use. Further, they have good potential for fishery. Thus small rural ponds can be a very good source of water for drinking, domestic use and also for generating income from fishery which can be augmented with scientific management as small reservoirs are more manageable and high yielding than larger ones. Hence, it is necessary to protect and conserve these small water bodies. This demands immediate action from ecologists, planners and policy makers.

REFERENCES

1. Ali S, Shaha S. and Mahmud N. (1985): Studies on the physico-chemical aspects of Maheshkhali channel, Bay of Bengal. Dhaka University Studies, B, XXXIII(1): 43-49.
2. Banerjee V. (1998): Influence of zinc and mercury on blood parameters of the fish *Heteropneustes fossilis*. Environ. Ecol., 16: 79-84.
3. Cairns J. and Dickson K.L. (1971): A simple method for the biological assessment of the effect of waste discharge on aquatic bottom dwelling organism. J. Water Pollut. Control Fed., 43: 122-275.
4. Choudhury P.R., Pandey R.A. and Bal A.S. (1998): Macrophyte infestation of water bodies and methods of lake restoration. In: Conservation and management of aquatic resources (Ed.: A. Gautam). Daya Publishing House, Delhi. pp. 200-224.
5. Das A.K. (2002): Limno-chemistry and productivity of upper Ganga complex. Poll. Res., 21: 157-168.
6. Dutta S, S Gupta and A Gupta (2004): Euglenoid blooms in the flood plain wetlands of Barak Valley, Assam, North-Eastern India. J. Environ Biol., 25: 369-373.
7. Gobler C.J. and Coper E.M. (1996): Stimulation of brown tide bloom by iron. In: Harmful and toxic algal bloom (Eds.: T. Yasumoto, Y. Oshima and Y. Fukuyo). Intergovt. Oceanogr. Commission, UNESCO, Paris. pp. 321-324.

8. Goldman C.R. (1960): Primary productivity and limiting factors in three lakes of the Alaska Peninsula. *Ecol. Monogr.*, 30: 207-230.
9. Gupta A. (1996). Heavy metals in water, periphytonic algae, detritus and insects from two streams in Shillong, Northeastern India. *Environ. Monit. Assess.*, 40: 215-223.
10. Gupta A.K. and Chakraborty P. (1995): Effect of zinc on the testes of *Notopterus notopterus* and its subsequent recovery by EDTA. *J. Inland Fish. Soc. India*, 27: 57-59.
11. Hasan G.O., Paul P. Mathisen and Don Pellegrino (2007): Distribution of heavy metals in vegetation surrounding the Blackstone River, USA: Considerations regarding sediment contamination and long term metals transport in freshwater riverine ecosystems. *J. Environ. Biol.*, 28: 493-502.
12. Khan M.A. and Bhatt G.H. (2000): Biological invasion and 'Red water' phenomenon in lake. Manasbal of Kashmir Valley, India. *Poll. Res.*, 19: 113-117.
13. Manna R.K. and Das A.K. (1984): Impact of the river moosi on river Krishna 1. *Limno-chemistry. Poll. Res.*, 23: 117-124.
14. Mitra S.K. and Bhattacharya B.K. (1965): Some hydrological factors affecting plankton production in fish pond in Kalyani, West Bengal, India. *Ichthyological*, 4: 8-12.
15. Pandit A.K. (2002): Algae as a component of Dal lake ecosystem in Kashmir Himalaya. In: *Ecology and conservation of lakes, reservoirs and rivers* (Ed.: Arvind Kumar). A.P.H. Publishing Corporation, New Delhi.
16. Park Bae Kyung and Seok Soon Park (2005): Effects of stream hydraulic conditions on foraging strategies of false dace, *Pseudorasbora parva* in the lentic ecosystem. *J. Environ. Biol.*, 26: 635-643.
17. Ramesh R. and Anbu M. (1996): *Chemical methods for environmental analysis: Water and sediment*. Macmillan India Ltd., Madras. p. 161.
18. Sharma L.L. and Sarang N. (2004): Physico-chemical limnology and productivity of Jaisamand Lake, Udaipur (Rajasthan). *Poll. Res.*, 23: 87-92.
19. Tiwari A. and Chauhan S.V.S. (2006): Seasonal phytoplankton diversity of Kitham Lake, Agra. *J. Environ. Biol.*, 27: 35-38.
20. Usha R., Ramalingam K. and Bharathi Ramjam U.D. (2006): Freshwater lakes- A potential source for aquaculture activities- A model study on Perumal Lake, Cuddalore, Tamil Nadu. *J. Environ. Biol.*, 27: 713-722.
21. Verma S.K. (2002): Fresh water toxic blue-green algal blooms - A response to extra nutrient enrichment. In: *Ecology of polluted waters. Voll. II* (Ed.: A. Kumar). A.P.H. Publishing Corporation. New Delhi. pp. 1161-1175.
22. Wetzel R.G. (1984): *Limnology*, Saunders College Publishing, New York. pp. 760.